

# STOP

## NSAIDS! and SMOKING!

The use of non-steroidal anti-inflammatory drugs (NSAIDS) and smoking are both associated with a **HIGH RISK OF DEVELOPING STOMACH ULCERS**

The following drugs are NSAIDs or include NSAIDS in their formula and should not be taken after weight loss surgery:

Advil	Clinoril	Indo-Lemmon	Orudis
Aleve	Combunox	Indomethagan	Oruvail
Amigesic	Darvon	Lodine	Pamprin IB
Anacin	Daypro	Lodine XL	Percodan
Anaprox	Disalcid	Meclomen	Ponstel
Anaprox DS	Dolobid	Micrainin	Relafen
Ansaid	EC Naprosyn	Midol IB	Rexolate
Arthrotec	Ecedrin	Mobic	Tab-Profen
Ascriptin	Ecotrin	Motrin	Tandearil
Aspirin	Equagesic	Motrin IB	Tolectin
Azolid	Exedrin IB	Nalfon	Tolectin 600
Bextra	Feldene	Nalfon 200	Tolectin DS
Bufferin	Ibuprin	Naprapac	Toradol
Butazolidin	Ibuprofen	Naprelan	Uracel
Cataflam	Indocin	Naprosyn	Vicoprofen
Celebrex	Indocin SR	Nupin	Voltaren