

# Colonoscopy Prep Instructions: Polyethylene Glycol 3350 Generic (Golytely, Nulytely or Trilyte)

**IMPORTANT** - Please Read These Instructions at Least 2 Weeks Before Your Colonoscopy.

If you need to reschedule, cancel, or have any questions about your procedure please contact 954.689.5556 before 5 p.m. or 954.689.5000 after hours

## **Key Instructions:**

Your doctor has scheduled you for a colonoscopy. To have a successful colonoscopy your colon must be clear of any stool. Follow all of the instructions in this handout **EXACTLY** as they are written. If you do not follow the directions, your procedure may be canceled.

- **Do NOT** eat solid food the ENTIRE day before your colonoscopy.
- Buy your bowel preparation at least **5 days** before your colonoscopy.
- **Do NOT** mix the solution until the day before your colonoscopy.
- A responsible family member or friend **MUST** drive you home from the procedure. You are **NOT ALLOWED** to drive, take a taxi, bus or car service, or leave the Endoscopy Center **ALONE**. If you do not have a responsible driver (family member or friend) with you to take you home, your exam cannot be done with sedation and may be canceled. You **MUST** remain in the endoscopy center until you recover from sedation. The entire process from arrival to discharge may take 3-4 hours.
- Please do not bring any jewelry or other valuable items to the endoscopy room for your procedure.
- Please bring a list of known allergies, current medications, a list of medical conditions and surgical history.

## **Medications**

### **The day before your colonoscopy:**

Some of the medicines you take may need to be stopped or adjusted temporarily before your colonoscopy.

- **Blood thinners** -- such as Coumadin® (warfarin), Plavix® (clopidogrel), Ticlid® (ticlopidine hydrochloride), Agrylin® (anagrelide), Xarelto® (Rivaroxaban), Pradaxa® (Dabigatran), Eliquis® (Apixaban), and Effient® (Prasugrel). You must speak with your prescribing physician or the specialist performing the procedure at least 2 weeks before the scheduled test. Do not stop these medications without the consent of your physician.

### **Insulin and diabetes medications**

- Please call the doctor that monitors your glucose levels. Your medications may need to be adjusted due to the diet restrictions required with this bowel preparation. The following is some information to guide you:
- If you take a pill to lower your sugar, do not take it on the day of your procedure.
- If you are taking regular insulin®, do not take it on the day of your procedure.
- If you are taking long-acting insulin (NPH), take half of your prescribed dose on the morning of the procedure.
- If you are taking any other insulin preparation, such as Lantus®, Humalog® or 70/30 insulin, contact your prescribing physician for instructions.
- If you take aspirin, take it and **ALL** other prescribed medicines with a sip of water on **the day of your colonoscopy**.

*Continued on back*

# Colonoscopy Prep Instructions: Polyethylene Glycol 3350 Generic (Golytely, Nulytely or Trilyte)

Continued from front

## FIVE (5) DAYS BEFORE YOUR COLONOSCOPY

- **Do NOT** take products that contain iron -- such as multi-vitamins -- (the label lists what is in the products).

Buy the prescription bowel preparation solution at your local pharmacy.

## THREE (3) DAYS BEFORE YOUR COLONOSCOPY

Avoid high-fiber foods -- such as popcorn, beans, seeds (flax, sunflower, and quinoa), multigrain bread, nuts, salad/vegetables, fresh and dried fruit. Avoid fiber supplements -- such as Metamucil®, Citrucel®, or Konsyl®.

## ONE (1) DAY BEFORE YOUR COLONOSCOPY

**Only drink clear liquids the ENTIRE DAY before your colonoscopy. Do NOT** eat any solid foods. The **clear liquids** you can drink include:

- Water, apple, or white grape juice; fat free broth; coffee or tea (without milk or creamer); clear carbonated beverages such as ginger ale or lemon-lime soda; Gatorade® or other sports drinks (not red); Kool-Aid® or other flavored drinks (not red). You may eat plain jello or other gelatins (not red) or popsicles (not red).

**Do NOT** drink alcohol on the day before or the day of the procedure.

## When to Drink Your Bowel Prep

Follow the instructions on the label. After mixing, place the solution in the refrigerator for a couple of hours before drinking. You may add the flavor packet that came with the bowel preparation. **DO NOT** add ice, sugar or any other flavorings to the solution.

## Morning Appointments

(Before 12 noon)

- OPTION 1 - Split dosing Prep** (*split dose is considered best option*)

### Step 1.

- Start drinking the bowel preparation at 6 PM the **evening before** your colonoscopy. Drink an 8-oz glass of bowel preparation every 10 minutes for a total of 12 glasses (3 liters).
- You may continue to drink clear liquids only until midnight.

### Step 2. The day of the colonoscopy (4 hours before your scheduled exam).

- Drink an 8-oz glass of bowel preparation every 10 minutes for a total of 4 glasses (1 liter).

**Please do not consume anything else by mouth after the bowel preparation is completed.**

- OPTION 2 - EVENING Before Preparation**

Start drinking the bowel preparation at **6 PM** the **evening before** your colonoscopy. Drink an 8-oz glass of bowel preparation every 10 minutes. You must finish drinking the solution by **9 PM** the night before your scheduled procedure. You can continue to drink clear liquids until midnight

**Please do not consume anything by mouth after midnight**

## Afternoon Appointments

(After 12 noon)

- Start drinking the bowel preparation at 6 AM the day of your colonoscopy. Drink an 8-oz glass of bowel preparation every 10 minutes. You must finish drinking the solution by 10 AM.
- You may continue to drink clear liquids up to 3 hours before your exam.