MEDICATION GUIDELINES BEFORE VASCULAR SURGERY

You MAY need to stop taking some medications before your surgery.

Please tell your doctor about EVERY medication you take. This includes prescription and nonprescription (over the counter items like aspirin, vitamins, herbs, etc.).

Carefully read the information below. Call your doctor or Surgical Coordinator if you have ANY questions.

DO NOT STOP TAKING ANY MEDICATION UNLESS YOUR DOCTOR OR NURSE TELLS YOU TO!

Follow the instructions for all CHECKED medications below:

Aspirin	
- Keep taking	
 Stop taking before surgery. Take your last dose on 	
Plavix (clopidogrel)	
- Keep taking	
 Stop taking before surgery. Take your last dose on_ 	
Coumadin (warfarin) - Stop taking before surgery. Ta	ke your last dose on
Lovenox - Take mg twice a day at 7am and 7pm	
Take your last dose on	after your 7am or 7pm injection.
Pletal (cilostazol) - Stop taking before surgery. Take y	your last dose on
Brilinta (ticagrelor) - Stop taking before surgery. Take	e your last dose on
Eliquis (apixaban) - Stop taking before surgery. Take	e your last dose on
Effient (prasugrel) - Stop taking before surgery. Take	your last dose on
Xarelto (rivaroxaban) - Stop taking before surgery. Ta	ake your last dose on

OTHER MEDICATIONS AND SUPPLEMENTS I TAKE
Due some instanctions for these and the time
Pre-surgery instructions for these medications:



www.clevelandclinic.org/heart