Attending to Visitor Concern's Regarding Visitation Expectations

BEFORE beginning to converse with your patient's visitor be mindful of YOUR OWN EMOTIONS. It is very easy for one's own emotions to get in the way of responding calmly, tenderly and empathically.

Listed below are a few additional items to consider when holding crucial conversations with visitors who are anxious, dread-filled or aggressive.

- 1. Use positive language when making requests.
- 2. Make requests in clear, positive and concrete language.

3. Be aware that when visitors hear a demand from us, they often see two options: to submit or rebel.

4. Express appreciation when your patient's visitor complies with your request.

5. Empathize with the visitor who is expressing reluctance.¹

The suggested responses are based upon the R.E.D.E to Communicate[®]: Foundations of Healthcare Communication course developed by caregivers at Cleveland Clinic.² There are three phases to the model. Phase One, known as *the Establishment of a Relationship* is recommended.

The mnemonic known as **S.A.V.E.**, is offered as guidance for communicating the recommendations and addressing the mismatched expectations visitors may be expressing. The mnemonic does not need to be followed in order. Below are a few samples of S.A.V.E. statements:

• S = SUPPORT

"I am here for you and your loved one (wife, husband, mother, father, etc) "I want to help you."

• A = ACKNOWLEDGE

"I can truly appreciate the worry you have about leaving your mother." "It's harder to manage our own worry when we aren't at his/her bedside." "I wish you and your loved one didn't have to go through any of this."

• V = VALIDATE

"You have every right to feel 'worried' 'troubled' 'anxious' 'frightened' 'unsettled' ... we are living in a very unpredictable time."

• **E = EMOTION NAMING** Identify the emotion the family member is conveying "I hear worry in your voice." "I am wondering if you are feeling overwhelmed."



Below is a sample script. Say only what comes natural to you. When we can empathize with the experiences of patients and families, we are able to connect with them in powerful ways. ³

SCENARIO 1. Too many visitors

Nurse: (Directed to patient) You certainly are popular! Please introduce me to your visitors.

Wait for response to identify the relationships of visitors to your patient Direct the following statements to the visitors in the room. Use the names of the visitors in the remainder of the conversation.

Nurse: <Visitor's name, > I am here to take the best care possible of <insert patient's name.> To do that I must enforce the new visitation policy that limits each patient to one visitor at a time. Were you aware of the new policy?

Wait for response ...

Nurse: I can imagine that this new rule has left you all unsettled as you all want to be with him/her. For <insert patient's name> safety and well-being, I must ask you all to decide upon who will be the one to remain with <patient's name.>. I will give you a minute to say good- bye and return in a few minutes to check on you, <patient's name.>

Return to the room in 10 minutes.

Nurse to remaining visitor: Thank you for accommodating my request. I know this was difficult for everyone. How are you doing?

Listen for patient's and visitor's response.

Nurse: That is helpful for me to know. I will share this important information with the rest of the team caring for you.

Nurse: Before I leave, I want to make sure that you know the new visitation policy so you can share it with the members of your family. We are limiting visitors to one person who is over the age of 16 between the hours of 8 am – 12 noon and 4 pm – 8 pm. It would be terrible for another family member to drive to the hospital and not be able to visit. In this unpredictable time, it is important that we all look out for one another. Thank you for your understanding.

SCENARIO 2. Visitors who will not leave

Nurse: It is 7:45 pm and I want to let you know that visiting hours will be ending in fifteen minutes.

If visitor says they won't leave, ask them to voice their concerns about leaving.

Nurse: Tell me about your concerns.

Listen to visitor's concerns. Allow them to express everything that worries them. Exhaust their concerns by asking, "Are there other concerns you have about leaving?"

Nurse: I hear the worry in your voice and I am grateful that you shared these concerns with me.

Nurse: I want you to know you are not alone in these worries. This is a very difficult time. No one wants to be away from their loved one, most especially when they're sick.

Nurse: <Use visitor's name,> I need to ask you to entrust your <dad's, mom's, etc.> care to me. Your Patient Care Nursing Assistant or PCNA, <insert name,> the nurses, and I will be looking after him/her while you are gone. The information you shared will help us take even better care of your <dad, mom, etc.> as we know his/her worries as well as yours.

Nurse: I want you to go home now and rest as best you can so you can continue to provide support to <patient's name>. If you wake up in the middle of the night and are worried, call the nursing station and ask to speak with your <dad's, mom's, etc.> nurse. I'll be happy to give you the number to call so you'll have it if you need it. In addition, we have your information and will call you if necessary. How does that sound to you?

Nurse: I am grateful for your understanding.

Escalating and Intense Emotion

Visitor refuses to leave. Listen to their reason(s) and pay particular attention to their emotion.

Nurse: I hear your intention and need to stay with your dad. I wish I could honor that need. I'm so sorry that this is difficult for you. If my dad were in the hospital, I would want to be at his bedside 24/7 too. (Pause. Wait for their response.)

Nurse: I am asking you to embrace the new visiting hours that have been established to keep you, your <dad, mom, etc.> safe, and all the people working in this hospital safe.

Nurse: Thank you for honoring my request.



*The goal with managing intense emotion is two-fold. First, remember that when we sense conflict, we tend to speak faster, louder, and with a higher pitch. To combat, take long slow deep breaths to slow our body's response. Second, listen fully to what is said and reflect back what you are hearing. Include empathy statements in response to the emotion conveyed. Reflective listening with empathy will slow their body's response.

SCENARIO 3: Unauthorized use of Hospital PPE by Visitors

Nurse: You want to wear protective items to keep your loved one and everyone else in the hospital safe. I'm grateful for your vigilance.

Nurse: I'm sorry that we can't provide these items for you. I wish we had unlimited resources. We must prioritize use of protective gear for individuals who are designated COVID 19 positive and their caregivers.

Nurse: The CDC has determined our best course of action for your loved one is to limit the number of visitors and establish a safe distance between <patient's name> and others. We've implemented those recommendations and ask that you observe them.

Nurse: Thank you for supporting us and our efforts to provide care for you and your loved one.

It can be helpful to have a reminder of the vocabulary of emotions and feelings patients, families and caregivers can experience during demanding, unpredictable and stressful times.

Feelings when needs are not fulfilled ¹					
Angry	Discouraged	Hopeless	Overwhelmed		
Annoyed	Disappointed	Impatient	Puzzled		
Concerned	Embarrassed	Irritated	Reluctant		
Confused	Frustrated	Lonely	Sad		
Disappointed	Helpless	Nervous	Uncomfortable		
		ANGER ⁴			
Soft Anger					
Annoyed	Crabby	Detached	Indifferent		
Apathetic	Cranky	Displeased	Irritated		
Bold	Critical	Frustrated	Peeved		
Cold	Cross	Impatient	Rankled		
Mood-State Anger		· · · · · · · · · · · · · · · · · · ·			
Affronted	Arrogant	Indignant	Resentful		
Aggravated	Bristling	Inflamed	Riled Up		
Angry	Exasperated	Mad	Sarcastic		
Antagonized	Incensed	Offended			
Intense Anger		·			
Aggressive	Disgusted	Livid	Seething		
Appalled	Furious	Outraged	Spiteful		
Belligerent	Hateful	Menacing	Vengeful		
Bitter	Hostile	Raging	Vicious		
Contemptuous	Irate	Ranting	Violent		
		FEAR ⁴			
Soft Fear					
Alert	Disconcerted	Indecisive	Timid		
Apprehensive	Disoriented	Insecure	Uneasy		
Cautious	Disquieted	Hesitant	Watchful		
Concerned	Edgy	Leery			
Confused	Fidgety	Pensive			
Curious	Hesitant	Shy			
Mood – State Fear					
Afraid	Fearful	Shaky	Wary		
Alarmed	Jumpy	Startled	Worried		
Anxious	Nervous	Suspicious			
Aversive	Perturbed	Unnerved			
Distrustful	Rattled	Unsettled			
Intense Fear					
Dread-filled	Panicked	Petrified	Terrified		
Horrified	Paralyzed	Shocked			

Here are some examples:

Authored by: Mary Beth Modic, DNP, RN, CNS; Edited by Lauren Peeples, BS and Carol Santalucia, MBA of the Center for Excellence in Healthcare Communication and 50 Health Care Partners (aka Patient Volunteers); Approved by Corporate Communications Version Date: 3/18/20 ©2020 THE CLEVELAND CLINIC FOUNDATION. ALL RIGHTS RESERVED.

Cleveland Clinic

Office of Patient Experience patientexperience@ccf.org

SADNESS ⁴					
Soft Sadness					
Disappointed	Distracted	Low	Wistful		
Disconnected	Listless	Regretful			
Mood – State Sadness					
Discouraged	Dispirited	Grieving	Sad		
Down	Drained	Heavyhearted	Sorrowful		
Downtrodden	Forlorn	Melancholy	Weepy		
Discouraged	Gloomy	Mournful	World-weary		
Intense Sadness					
Anguished	Despairing	Grief-stricken	Inconsolable		
Bleak	Despondent	Heart-broken	Morose		
Bereaved	Depressed	Hopeless			

References:

1. Rosenberg, M.B. 2015. *Nonviolent Communication: A Language of Life*. 3rd ed. Encinitas, CA.: PuddleDancer

2. Windover, A.K., Boissy, A., Rice, T.W., Gilligan, T., Velez, V.J & Merlino, J. (2014). The REDE Model of Communication: Optimizing Relationship as a Therapeutic Agent. *Journal of Patient Experience*, *1* (1), 8-13.

3. Koloroutis, M. & Trout, M. (2012). See me as a person. Minneapolis: Creative Healthcare Management.

4. McLaren, K. (2012). *The Art of Empathy*. Boulder: Sounds True.

For additional messaging and resources, please visit <u>http://connect.ccf.org/CEHC</u>