

Recommendations For Personal Protective Equipment NON-CLINICAL CAREGIVERS of Patients COVID-19 Positive or Persons Under Investigation (PUI)

	Face Mask (surgical or ear-loop mask)	N95 (fit-tested respirator)	Protective Eyewear (goggles or faceshield)	Gown & Gloves
Caregivers Performing Screening at Point of Entry to Hospital or Department	Wear Face Mask *		Wear Protective Eyewear **	
Patient Service Representative (PSR) • if serving as point of entry screener follow above recommendation for “Caregivers Performing Screening”	Follow Standard Precautions			
Health Unit Coordinator (HUC)	Follow Standard Precautions			
Dietary/Nutrition	Advised not to enter room. Leave trays at nursing station			
Patient Transporters of patients in Contact/Droplet + Eyewear & Specified Precautions—UPON <u>ENTRY</u> INTO PATIENT ROOM	Wear Face Mask			Wear Gown & Gloves
Patient Transporters of patients in Contact/Droplet + Eyewear & Specified Precautions—UPON <u>EXITING</u> PATIENT ROOM	Patient and Transporter: Wear <u>new</u> Face Mask Upon exiting			Only if direct contact with patient or contact with contaminated equipment
EVS doing Routine Cleaning for rooms with Contact/Droplet Isolation	Wear Face Mask		Wear Protective Eyewear	Wear Gown & Gloves
EVS doing Terminal (Discharge) Cleaning for rooms with Contact/Droplet Isolation	Wear Face Mask		Wear Protective Eyewear (if splashes or sprays anticipated)	Wear Gown & Gloves
Bus/Shuttle Drivers/Valet	Wear Face Mask *			
Materials Handlers	Follow Standard Precautions			

***Face mask usage:** Extended use for face masks: One mask per shift. Change face mask with contamination with respiratory secretions, or obvious soilage or damage. Masks should always be worn covering nose & mouth. Used mask handled by ear loops ONLY. Never touch the front of the mask. Perform hand hygiene after handling used mask. Do not reapply a used mask.

****Protective Eyewear:** Reusable goggles or facemask preferable. Can be worn for extended period. Must be cleaned with disinfectant wipes with contamination/soilage and after each use.

Direct contact :

Distance: within 6 feet of patient
 Time: >10 minutes
 Contact with patient’s secretions, body fluids, blood, etc.

Standard Precautions

- Hand hygiene
 Cough etiquette
 Personal Protective Equipment (PPE)
- Gloves: if reasonably expect to come into contact with non-intact skin or body fluids
 - Gowns: if reasonably expect that your uniform could get soiled
 - Masks / Face protection: during care activities where this a chance of splash or spray of body fluids, blood, excretions, secretions